ลิงลม

ຕັກນ໙ະ:	กวามยาวลำตัวและหัว 30-38 ซม. หางยาว 1-2 ซม. น้ำหนัก 1-2 กิโลกรัม เป็นสัตว์ใน กลุ่มลิงที่มีขนาดเล็ก รูปร่างอ้วนกลม หัวกลม ตากลม รอบดวงตาเป็นวงสีเข้ม หูเล็กและ
	สั้น ขนสั้นหนานุ่มสีน้ำตาลหรือสีน้ำตาลแกมเทา ค้านท้องสีอ่อนกว่า ตลอดแนวของสัน หลังมีแถบสีคำเห็นเด่นชัด หางสั้น ตัวเมียมีเต้านม 2 คู่
เขตแพร่กระจาย:	หแงมแบบแท แทนเทน ขุด กางเาน ๆ มมอมเตานม 2 กู เมียนมาร์ ไทย ลาว กัมพูชา เวียดนาม มาเลเชีย เกาะสุมาตราและเกาะบอร์เนียวของ อิน
	โคนีเชีย และเกาะมินคาเนาของฟิลิปปินส์
ที่อยู่อาศัย:	อาศัยอยู่ได้ในป่าหลายประเภท ทั้งป่าสมบูรณ์และป่าเสื่อมโทรม ป่าสะแกราชพบในป่า
च् च	ดิบแล้ง เช่น บริเวณ กม.4
อุปนิสัย:	โดยทั่วไปจะเห็นอยู่โดดเดี่ยว ออกหากินเวลากลางคืน กลางวันนอนหลับพักผ่อนบนคา
	ุ คบไม้หรือตามพุ่มไม้ใบหนา ปกติจะไม่ลงมายังพื้นดิน เคลื่อนไหวเชื่องช้า แต่จะว่องไว มากเมื่อพบเหยื่อ อาหารได้แก่ แมลง นก สัตว์ขนาดเล็ก และยังกินลูกไม้สุก น้ำหวานและ
	เกสรคอกไม้ ตกลูกครั้งละ 1 ตัว ระยะตั้งท้อง 90 วัน ให้นมลูกประมาณ 190 วัน ลูกจะอยู่
	กับแม่ตลอคระยะ 6-9 เคือนแรก
สถานภาพ:	พบไม่บ่อยนัก และกฎหมายจัดเป็นสัตว์ป่ากุ้มกรอง
Characteristics:	Head and body 30-38 cm; tail 1-2 cm; weight 1-2 kg. A small-sized primate with plump
	body and rounded head. Its large rounded eyes are surrounded by dark rings; ears small
	and short; body covered with thick soft fur of brownish or grayish brown color,
	distinctly furnished along the spinal line with a dark band; belly paler in color; tail
	remarkably short. A female has 2 pairs of mammae.
Distribution:	Myanmar, Thailand, Laos, Vietnam, Kampuchea, Malaysia, to Sumatra and Borneo of
	Indonesia, and Mindanao island of the Philippines.
Habitat:	Frequents in various types of forest, both the healthy and deteriolated ones. In Sakaerat
	area, it lives in dry evergreen forest, notably at the km 4.
Habit:	Generally seen as a solitary animal. Foraging in the night time, and rests in tree forks or
	hides among thick foliage in the day time. It rarely descends to the ground, but normally
	moves with awkward gait between branches. On capturing preys it can do with

surprisingly great speeds; which consisting of insects, birds, small animals, and also ripe fruits, nectar and pollen. One young is borned at a time, after a gestation period of 90 days. Young stays with its mother for the first 6-9 months during which its mother suckles it for 190 days.

Status:

Seldomly seen, and being protected legally as a protected wild animal.





ลิงลม (ชื่ออื่น: นางอาย)

Common Slow Loris

Nycticebus coucang (Boddaert, 1785)